

Constitutively Active Receptors

Figure 1 (Page 1 of 15)

CLASS A GROUP II						
A1AB_human	α_{1B} -adrenergic alpha 1B-AR	TM ^{DI}	⁶³ FAIVGNILVIL A	IP / CCS-7	(Scheer, Fanelli et al. 1997)	
		junction between TM ^{DI} and IC ²	¹⁴² CAISIDRYIGV A			
A1AB_human	α_{1B} -adrenergic alpha 1B-AR	junction between TM ^{DI} and IC ²	¹⁴³ CAISIDRYIGV K	IP / CCS-7	(Scheer, Costa et al. 2000)	
A1AB_human	α_{1B} -adrenergic alpha 1B-AR	TM ^{III}	¹²⁸ AVDV _{LC} CTASII F	IP / CCS-1	(Perez, Hwa et al. 1996)	
		carboxyl end of IC ³	²⁹³ REKK _{AA} AKTLGII E	IP arachidonic acid release		
		TM ^V	²⁰⁴ EFPFY _{AL} FSSLLG V	IP / CCS-1	(Hwa, Garwin et al. 1997)	
A1AB_human	α_{1B} -adrenergic	C-terminal IC ³	²⁹³ SREKK _{AA} AKT X=19 different substitutions	PI / CCS-7	(Kjelsberg, Cotecchia et al. 1992)	
A1AB_human	α_{1B} -adrenergic	C-terminus IC ³	²⁸⁸ KFSREKK _{AA} AKTLGII K H L	PI hydrolysis / rat fibroblast	(Allen, Lefkowitz et al. 1991)	
A2AA_human	α_2 C10-adrenergic alpha-2AAR	C-terminal IC ³ loop	³⁷³ (348?) EKRF T FVLAV X=F, A, C, E, K	adenylyl cyclase inhibition / HEK293	(Ren, Kurose et al. 1993)	
ACM1_human	muscarnic Hm1	C-terminal IC ³ loop junction	³⁶⁰ SLV KE KKAAARTLS A	PI / HEK293	(Högger, Shockley et al. 1995)	
ACM2-human	muscarnic acetylcholine M1 muscarnic acetylcholine M2	junction of IC ³ and TM ^{VI}	³⁹⁰ KKV PT TIL ¹ A 1-4 A inserted	IP production, inhibition of cAMP production / COS-7	(Liu, Blin et al. 1996)	

Figure 1 (Page 2 of 15)

CLASS A GROUP II						
ACM3_rat	m3 muscarinic (rat)	TMVI	507 TWTPYNNIMVLVNT S	IP / COS-7	(Bluml, Mutschler et al. 1994)	
ACM5_human	muscarinic acetylcholine M3 m5 muscarinic	N-terminus to TMII	chimera composed of m2 1-69 m5 77-445 m2 391-466	β-gal / NIH 3T3	(Burstein, Spalding et al. 1996)	
ACM5_human	m5 muscarinic muscarinic acetylcholine M5	TMVI	451 A _M LLA E _L ITW T _V PYNNI MVLVST V S T	β-gal; radioligand binding / NIH-3T3	(Spalding, Burstein et al. 1998)	
ACM5_human	m5 muscarinic muscarinic acetylcholine M5	junction of TMVI and EC3	465 YNIIMVLV _S TFCDFKCV X=V,F,R,K,+more	β-gal; radioligand binding / NIH-3T3	(Spalding, Burstein et al. 1997)	
B1AR_human	β ₁ -adrenergic	C-terminus	389 RKA _R QGLLCCA	adenylyl cyclase; agonist binding / CHW	(Mason, Moore et al. 1999)	
B2AR_human	β ₂ -adrenergic beta-2AR	C-terminal IC3 loop	266 FCLKEH _S K _R A _K KT _A LG _I	adenylyl cyclase activation; agonist binding affinity / COS-7 or CHO	(Samama, Cotecchia et al. 1993); (Lefkowitz, Cotecchia et al. 1993)	
DADR_human	dopamine D1A	carboxyl terminal IC3	264 SFKMSE _I K _R E _K T _V LKT 288 from D1B receptor	adenylyl cyclase; cAMP accumulation / HEK293	(Charpentier, Jarvie et al. 1996)	
DADR_human	dopamine D1	TMVI	286 APDTSIKK _A E _K T _V LKT FVCCW _L PPFFIL A	cAMP accumulation / COS-7	(Cho, Taylor et al. 1996)	
HH2R_rat	histamine H ₂	IC2	115 FMISL _D RYCAV N,A	cAMP production / HEK-293	(Alewijne, Timmerman et al. 2000)	

Figure 1 (Page 3 of 15)

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP III					
OPSD_human	opsin rhodopsin	TMII TMIII TMVII	90 FWVILGG G FTSTLY D 113 GCN L EGFFAT Q 292 296 MTIPAFFAKSAAIY E G, E, M ²⁹² Ala neutral a.a converted to carboxylate and competes with ¹³ Glu for salt bridge with ²⁹⁶ Lys	transducin; phosphorylation by rhodopsin kinase / COS	(Rim and Oprian 1995)
OPSD_human	opsin rhodopsin	TMIII	134 VVLAI E RYVVV I, Q, S	transducin; radioligand binding / COS	(Acharya and Karnik 1996)
OPSD_human	opsin rhodopsin	TM6 <i>plus</i> TM3 TMVII	257 RMVIIIMVIAFL Y, N <i>plus</i> G113Q 296 PAFFAKSAAIY G X=E, M natural mutants + 10 different a.a. substitutions	transducin, GTP γ S uptake / COS transducin; radioligand binding / COS	(Han, Smith et al. 1998) (Govardhan and Oprian 1994); (Cohen, Yang et al. 1993)
OPSD_human	opsin rhodopsin	IC2	134 VVLAI E RYVVV Q	disrupts critical salt bridge between ²⁹⁶ Lys(TMVII) and ¹³ Glu(TMIII)	(Cohen, Yang et al. 1993)

Figure 1 (Page 4 of 15)

TRFR_mouse	thyrotropin-releasing hormone TRH-R	carboxyl tail FRKLQNCKQK STOP	³³⁵ "Ca ²⁺ efflux, [Ca ²⁺] / Xenopus oocytes; IP formation / AtT20, <i>stably transfected</i>	(Matus-Leibovitch, Nusgenszveig et al. 1995)

Figure 1 (Page 5 of 15)

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A					
GROUP IV					
BRB2_human	bradykinin B ₂	TMIII	AIISM ^N LYSSI	IP production / COS-7	(Marie, Koch et al. 1999)
	B2 bradykinin	TMVI	A ²⁵⁶		
	BK-2		LLF ^I IIC ^M LPFQI ^F		

Figure 1 (Page 6 of 15)

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP V AG2R_rat	AT _{1A} Type-1A angiotensin II	TMIII	111 ASV F NLYASV A disrupts ¹¹¹ Asn (TMIII) - ²⁹² Tyr (TMVII) interaction	phospholipase C; IP production / COS-7	(Groblewski, Maigret et al. 1997)
AG2R_rat	AT _{1A} Type-1A angiotensin II	C-terminus of TM7 other multiple mutations	305 LFY G LGKKFK Q	IP production / HEK- 293; intracellular Ca ²⁺ mobilization / CHO	(Parnot, Bardin et al. 2000)
FMLR_human	formylmethionylleucylphenylal anine (MLPR)	IC1	51 LVIWVAG F RM I HTV T ISYLNKAVA LvvWVTAPEAKRTINA I WFLNLAVA (K above conflicts with SWISS-PROT database)	PI production; phospholipase C stimulation / COS-7	(Amatruda, Draga- Graonic et al. 1995)
IL8_human	interleukin-8 receptor B CXCR-2 chemokine	IC2	138 ACISV D RYLAIVH V	IP production; Ca ²⁺ mobilization and actin polymerization / NIH 3T3	(Burger, Burger et al. 1999)
LSHR_human	luteinizing hormone (LH)	IC3	564 MATNK D TKIAKK G	cAMP production / HEK293	(Kudo, Osuga et al. 1996)
LSHR_human	luteinizing hormone (LH)	TMVI	578 ILLIFT D FTCMA G	cAMP production / COS-7	(Shenker, Laue et al. 1993)
LSHR_human	luteinizing hormone (LH)	TM6	571 KIAKK M AILIFT D FTCM I I	cAMP production / COS-7	(Kosugi, Van Dop et al. 1995)
LSHR_rat	luteinizing hormone / human chorionic gonadotropin (LH/hCG)	TMVI	556 ILLIFT D FTCMA G, Y	cAMP production / HEK 293T	(Bradbury, Kawate et al. 1997; Bradbury and Menon 1999)
OPRD_mouse	delta opioid receptor	TM3	128 KV L SD D YYNM F A, K, H	adenylyl cyclase inhibition / COS-7	(Cavalli, Babey et al. 1999)
OXYR_human	oxytocin	IC2	137 LMSLD R CLAI C A	IP production / COS-7	(Fanelli, Barbier et al. 1999)

Figure 1 (Page 7 of 15)

PAFR_human	platelet-activating factor (PAF)	C-terminus of IC3	<u>EVKRRALWMVCTVLAV</u> R	IP production / CCS-7	(Parent, Le Gouill et al. 1996)
PAFR_human	platelet-activating factor (PAF)	TMIII	<u>CLFFINTYCSV</u> A	IP production, adenylyl cyclase inhibition / CHO	(Ishii, Izumi et al. 1997)
PE23_human	prostaglandin E ₃ , EP3II EP3IV	C-terminal tail	<u>FCQQEEFWGN</u> FCQMRKRLREQQEEFWGN	inhibition of adenylyl cyclase / CHO-K1	(Jin, Mao et al. 1997)
PE23_mouse	prostaglandin E ₃ , EP3	carboxyl-terminal tail	<u>↑truncated</u> 336 KILLRKFCQ <u>IRDEH</u> MMNHL <u>↑truncated</u> (3 α) (3 β)	inhibition of adenylyl cyclase / CHO, <i>stably expressed</i>	(Hasegawa, Negishi et al. 1996)
THTR_human	thrombin	EC2 loop	<u>CHDVLN<u>NET</u>LL<u>EGYYAYY</u></u> DLKD KDF I	⁴⁵ Ca ²⁺ efflux, PI hydrolysis, reporter gene induction / COS-7	(Nanovicz, Wang et al. 1996)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	EC1 EC2	<u>YYNN<u>HAIDWQTG</u></u> F, M 568 YAKV <u>SICL</u> PM <u>D</u>	inositol phosphate-diacylglycerol cascade / COS-7	(Parma, Van Sande et al. 1995)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	TMIII TMVII	<u>ASE<u>LSVY</u>YTLTV</u> A 672 YPL <u>NSCANPFL</u>	adenylyl cyclase activation / COS-7	(Duprez, Parma et al. 1994)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	TMV	<u>VAFV<u>IV</u>CCCHV</u> L	cAMP formation / COS-7 cells	(Esapa, Duprez et al. 1999)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	TMVII	<u>CAN<u>PFLYAI</u>FT</u> V	cAMP formation / CHO cells	(Russo, Wong et al. 1999)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	IC3	<u>VRN<u>POYNPGDKDTKIAK</u></u> <u>deletion</u>	cAMP formation / COS-7	(Wonerow, Schoneberg et al. 1998)

Figure 1 (Page 8 of 15)

TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	IC3 / TMVI	623 KDTKIAKRM ^V AVLIFT ^I DFICM	cAMP activation / COS-7	(Paschke, Tonacchera et al. 1994)
V2R_human	vasopressin V2	IC2	136 LAMTLD ^A RHRAI	cAMP formation / COS-7	(Morin, Cotte et al. 1998)

Figure 1 (Page 9 of 15)

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS B GROUP I CALR_human hCTR-1	human calcitonin hCTR-1 hCTR-2	wild type (native) protein		adenylyl cyclase cAMP production / COS-1	(Cohen, Thaw et al. 1997)
CLASS B GROUP II PTRR_human	parathyroid hormone PTH / PTH-related peptide	junction of IC1 and TMII junction of IC3 and TMVI	223 TRNYI ^H MHLFL R, K 410 KLIKSTLVLM ^P C, others	cAMP accumulation / COS-7	(Schipani, Jensen et al. 1997)
CLASS B GROUP III GIPR_human	glucose-dependent insulinotropic peptide (GIP-R)	TMVI	340 VFAPV ^T EEQAR ^P	cAMP production / L293	(Tseng and Lin 1997)
GLR_rat	glucagon	junction of IC loop1 and TMII IC end of TMVI	178 TRNYI ^H GNLFA ^R 352 RLARSTLLIP ^A	cAMP accumulation / COS-7	(Hjorth, Orskov et al. 1998)
VIPR_human	vasoactive intestinal peptide 1 (VIP)	junction of IC loop 1 and TMII junction of IC loop 3 and TMVI	178 RNYI ^H MHLFI ^R 343 LARSTLLIP ^X X= K, P	cAMP production / COS-7 or CHO	(Gaudin, Maoret et al. 1998) (Gaudin, Rouyer-Fessard et al. 1998)

Figure 1 (Page 10 of 15)

Figure 1 (Page 11 of 15)

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS_D					
O74283	pheromone	TM6	229 PLSAVQIYLGTP	heterologous yeast assay	(Olesnick, Brown et al. 1999)
RCB2					
C. cinereus	pheromone α -factor	TM6	258 QSLLVPSIIIFI	<i>lacZ</i> reporter gene	(Konopka, Margarit et al. 1996)
STE2_yeast	pheromone α -factor	double mutations TM5 and TM6	223 MSFVLYVVK N ILAIR C C 247 251 DSFPHILL N EQSLL CC CC double mutations double mutations	<i>lacZ</i> reporter gene / yeast	(Dube, DeCostanzo et al. 2000)
STE3_yeast	pheromone α -factor	IC3	194 DVRDILHCTNS Q	β -galactosidase	(Boone, Davis et al. 1993)
STE2_yeast	pheromone α -factor	TM6	253 258 LIMSCQSLLVPSIIIFI L LP	β -galactosidase	(Sommers, Martin et al. 2000)

Figure 1 (Page 12 of 15)

Bibliography

Acharya, S. and S. S. Kamik (1996). "Modulation of GDP release from transducin by the conserved Glu134-Arg135 sequence in rhodopsin." *J. Biol. Chem.* **271**(41): 25406-11.

Alewijne, A. E. H. Timmerman, et al. (2000). "The Effect of Mutations in the DRY Motif on the Constitutive Activity and Structural Instability of the Histamine H(2) Receptor." *Mol Pharmacol* **57**(5): 890-898.

Allen, L. F., R. J. Lefkowitz, et al. (1991). "G-protein-coupled receptor genes as protooncogenes: constitutively activating mutation of the alpha 1B-adrenergic receptor enhances mitogenesis and tumorigenicity." *Proc. Natl. Acad. Sci. U.S.A.* **88**(24): 11354-8.

Amaralda, T. T., 3rd, S. Dragas-Graonic, et al. (1995). "Signal transduction by the formyl peptide receptor. Studies using chimeric receptors and site-directed mutagenesis define a novel domain for interaction with G-proteins." *J. Biol. Chem.* **270**(47): 28010-3.

Blüml, K., E. Mutzschler, et al. (1994). "Functional role in ligand binding and receptor activation of an asparagine residue present in the sixth transmembrane domain of all muscarinic acetylcholine receptors." *J. Biol. Chem.* **269**(29): 18870-6.

Boone, C., N. G. Davis, et al. (1993). "Mutations that alter the third cytoplasmic loop of the α -factor receptor lead to a constitutive and hypersensitive phenotype." *Proc. Natl. Acad. Sci. U.S.A.* **90**(21): 9921-5.

Bradbury, F. A., N. Kawate, et al. (1997). "Post-translational processing in the Golgi plays a critical role in the trafficking of the luteinizing hormone/human chorionic gonadotropin receptor to the cell surface." *J. Biol. Chem.* **272**(9): 5921-6.

Bradbury, F. A. and K. M. Menon (1999). "Evidence that constitutively active luteinizing hormone/human chorionic gonadotropin receptors are rapidly internalized." *Biochemistry* **38**(27): 8703-12.

Burger, M., J. A. Burger, et al. (1999). "Point mutation causing constitutive signaling of CXCR2 leads to transforming activity similar to Kaposi's sarcoma herpesvirus-G protein-coupled receptor." *J. Immunol.* **163**(4): 2017-22.

Burstein, E. S., T. A. Spalding, et al. (1996). "Constitutive activation of chimeric m2/m5 muscarinic receptors and delineation of G-protein coupling selectivity domains." *Biochem. Pharmacol.* **51**(4): 539-44.

Cavalli, A., A. M. Babey, et al. (1999). "Altered adenylyl cyclase responsiveness subsequent to point mutations of Asp 128 in the third transmembrane domain of the delta-opioid receptor." *Neuroscience* **93**(3): 1025-31.

Champertier, S., K. R. Jarvie, et al. (1996). "Silencing of the constitutive activity of the dopamine D1B receptor. Reciprocal mutations between D1 receptor subtypes delineate residues underlying activation properties." *J. Biol. Chem.* **271**(45): 28071-6.

Cho, W., L. P. Taylor, et al. (1996). "Mutagenesis of residues adjacent to transmembrane prolines alters D1 dopamine receptor binding and signal transduction." *Mol. Pharmacol.* **50**(5): 1338-45.

Cohen, D. P., C. N. Thaw, et al. (1997). "Human calcitonin receptors exhibit agonist-independent (constitutive) signaling activity." *Endocrinology* **138**(4): 1400-5.

Cohen, G. B., T. Yang, et al. (1993). "Constitutive activation of opsin: influence of charge at position 134 and size at position 296." *Biochemistry* **32**(23): 6111-5.

Dube, P., A. DeCostanzo, et al. (2000). "Interaction between transmembrane domains five and six of the alpha -factor receptor." *J. Biol. Chem.* **275**(34): 26492-9.

Duprez, L., J. Parma, et al. (1994). "Germline mutations in the thyrotropin receptor gene cause non- autoimmune autosomal dominant hyperthyroidism." *Nat. Genet.* **7**(3): 396-401.

Egan, C. T., K. Herrick-Davis, et al. (1998). "Creation of a constitutively activated state of the 5-hydroxytryptamine2A receptor by site-directed mutagenesis: inverse agonist activity of antipsychotic drugs." *J. Pharmacol. Exp. Ther.* **286**(1): 85-90.

Esapa, C. T., L. Duprez, et al. (1999). "A novel thyrotropin receptor mutation in an infant with severe thyrotoxicosis." *Thyroid* **9**(10): 1005-10.

Fanelli, F., P. Barbier, et al. (1999). "Activation mechanism of human oxytocin receptor: a combined study of experimental and computer-simulated mutagenesis." *Mol. Pharmacol.* **56**(1): 214-25.

Gaudin, P., J. J. Maoret, et al. (1998). "Constitutive activation of the human vasoactive intestinal peptide 1 receptor, a member of the new class II family of G protein-coupled receptors." *J. Biol. Chem.* **273**(9): 4990-6.

Gaudin, P., C. Bouvier-Fessard, et al. (1998). "Constitutive activation of the human VIP1 receptor." *Ann. N.Y. Acad. Sci.* **865**: 382-5.

Govardhan, C. P. and D. D. Oprian (1994). "Active site-directed inactivation of constitutively active mutants of rhodopsin." *J Biol Chem* **269**(9): 6524-7.

Groblewski, T., B. Maigret, et al. (1997). "Mutation of Asn11 in the third transmembrane domain of the AT1A angiotensin II receptor induces its constitutive activation." *J Biol Chem* **272**(3): 1822-6.

Han, M., S. O. Smith, et al. (1998). "Constitutive activation of opsin by mutation of methionine 257 on transmembrane helix 6." *Biochemistry* **37**(22): 8253-61.

Hasegawa, H., M. Negishi, et al. (1996). "Two isoforms of the prostaglandin E receptor EP3 subtype different in agonist-independent constitutive activity." *J Biol Chem* **271**(4): 1857-60.

Herrick-Davis, K., C. Egan, et al. (1997). "Activating mutations of the serotonin 5-HT2C receptor." *J Neurochem* **69**(3): 1138-44.

Hjorth, S. A., C. Orskov, et al. (1998). "Constitutive activity of glucagon receptor mutants." *Mol Endocrinol* **12**(1): 78-86.

Högger, P., M. S. Shockley, et al. (1995). "Activating and inactivating mutations in N- and C-terminal i3 loop junctions of muscarinic acetylcholine Hm1 receptors." *J Biol Chem* **270**(13): 7405-10.

Hwa, J., R. Gaivin, et al. (1997). "Synergism of constitutive activity in alpha 1-adrenergic receptor activation." *Biochemistry* **36**(3): 633-9.

Ishii, I., T. Izumi, et al. (1997). "Alanine exchanges of polar amino acids in the transmembrane domains of a platelet-activating factor receptor generate both constitutively active and inactive mutants." *J Biol Chem* **272**(12): 7846-54.

Jensen, A. A., T. A. Spalding, et al. (2000). "Functional importance of the Ala116-Pro136 region in the calcium-sensing receptor. CONSTITUTIVE ACTIVITY AND INVERSE AGONISM IN A FAMILY CG-PROTEIN-COUPLED RECEPTOR [In Process Citation]." *J Biol Chem* **275**(38): 29547-55.

Jin, J., G. F. Mao, et al. (1997). "Constitutive activity of human prostaglandin E receptor EP3 isoforms." *British J Pharmacol* **121**: 317-23.

Krebsberg, M. A., S. Cotecchia, et al. (1992). "Constitutive activation of the alpha 1B-adrenergic receptor by all amino acid substitutions at a single site. Evidence for a region which constrains receptor activation." *J Biol Chem* **267**(3): 1430-3.

Konopka, J. B., S. M. Margarit, et al. (1996). "Mutation of Pro-238 in transmembrane domain 6 constitutively activates the G protein-coupled alpha-factor receptor." *Proc Natl Acad Sci U S A* **93**(13): 6764-9.

Kosugi, S., C. Van Dop, et al. (1995). "Characterization of heterogeneous mutations causing constitutive activation of the luteinizing hormone receptor in familial male precocious puberty." *Hum Mol Genet* **4**(2): 183-8.

Kudo, M., Y. Osuga, et al. (1996). "Transmembrane regions V and VI of the human luteinizing hormone receptor are required for constitutive activation by a mutation in the third intracellular loop." *J Biol Chem* **271**(37): 22470-8.

Lefkowitz, R. J., S. Cotecchia, et al. (1993). "Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins." *Trends Pharmacol Sci* **14**(8): 303-7.

Liu, J., N. Blin, et al. (1996). "Molecular mechanisms involved in muscarinic acetylcholine receptor-mediated G protein activation studied by insertion mutagenesis." *J Biol Chem* **271**(11): 6172-8.

Marie, J., C. Koch, et al. (1999). "Constitutive activation of the human bradykinin B2 receptor induced by mutations in transmembrane helices III and VI." *Mol Pharmacol* **55**(1): 92-101.

Mason, D. A., J. D. Moore, et al. (1999). "A gain-of-function polymorphism in a G-protein coupling domain of the human beta1-adrenergic receptor." *J Biol Chem* **274**(18): 12670-4.

Matus-Leibovitch, N., D. R. Nussenzveig, et al. (1995). "Truncation of the thyrotropin-releasing hormone receptor carboxyl tail causes constitutive activity and leads to impaired responsiveness in Xenopus oocytes and AtT20 cells." *J Biol Chem* **270**(3): 1041-7.

Morin, D., N. Cotte, et al. (1998). "The D136A mutation of the V2 vasopressin receptor induces a constitutive activity which permits discrimination between antagonists with partial agonist and inverse agonist activities." *FEBS Lett* **441**(3): 470-5.

Nanevici, T., L. Wang, et al. (1996). "Thrombin receptor activating mutations. Alteration of an extracellular agonist recognition domain causes constitutive signaling." *J Biol Chem* **271**(2): 702-6.

Olesnicky, N. S., A. J. Brown, et al. (1999). "A constitutively active G-protein-coupled receptor causes mating self-compatibility in the mushroom Coprinus." *Embo J* **18**(10): 2756-63.

Parent, J. L., C. Le Gouill, et al. (1996). "Mutations of two adjacent amino acids generate inactive and constitutively active forms of the human platelet-activating factor receptor." *J Biol Chem* **271**(14): 7949-55.

Figure 1 (Page 14 of 15)

Parma, J., J. Van Sande, et al. (1995). "Somatic mutations causing constitutive activity of the thyrotropin receptor are the major cause of hyperfunctioning thyroid adenomas: identification of additional mutations activating both the cyclic adenosine 3',5'-monophosphate and inositol phosphate-Ca2+ cascades." *Mol Endocrinol* 9(6): 725-33.

Parnot, C., S. Bardin, et al. (2000). "Systematic identification of mutations that constitutively activate the angiotensin II type 1A receptor by screening a randomly mutated cDNA library with an original pharmacological bioassay." *Proc Natl Acad Sci U S A* 97(13): 7615-20.

Paschke, R., M. Tonacchera, et al. (1994). "Identification and functional characterization of two new somatic mutations causing constitutive activation of the thyrotropin receptor in hyperfunctioning autonomous adenomas of the thyroid." *J Clin Endocrinol Metab* 79(6): 1785-9.

Pauwels, P. J., A. Gouble, et al. (1999). "Activation of constitutive 5-hydroxytryptamine 1B receptor by a series of mutations in the BXXXB motif: positioning of the third intracellular loop distal junction and its goalpha protein interactions [In Process Citation]." *Biochem J* 343 Pt 2: 435-42.

Perez, D. M., J. Hwa, et al. (1996). "Constitutive activation of a single effector pathway: evidence for multiple activation states of a G protein-coupled receptor." *Mol Pharmacol* 49(1): 112-22.

Ren, Q., H. Kurose, et al. (1993). "Constitutively active mutants of the alpha 2-adrenergic receptor [published erratum appears in J Biol Chem 1994 Jan 14;269(2):1566]." *J Biol Chem* 268(22): 16483-7.

Rim, J. and D. D. Oprian (1995). "Constitutive activation of opsin: interaction of mutants with rhodopsin kinase and arrestin." *Biochemistry* 34(37): 11938-45.

Robbins, L. S., J. H. Nadeau, et al. (1993). "Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function." *Cell* 72(6): 827-34.

Russo, D., M. G. Wong, et al. (1999). "A Val 677 activating mutation of the thyrotropin receptor in a Hurthle cell thyroid carcinoma associated with thyrotoxicosis." *Thyroid* 9(1): 13-7.

Samama, P., S. Cotecchia, et al. (1993). "A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model." *Journal of Biological Chemistry* 268(7): 4625-36.

Scheer, A., T. Costa, et al. (2000). "Mutational analysis of the highly conserved arginine within the Glu/Asp-Arg-Tyr motif of the alpha(1b)-adrenergic receptor: effects on receptor isomerization and activation." *Mol Pharmacol* 57(2): 219-31.

Scheer, A., F. Fanelli, et al. (1997). "The activation process of the alpha 1B-adrenergic receptor: potential role of protonation and hydrophobicity of a highly conserved aspartate." *Proc Natl Acad Sci U S A* 94(3): 808-13.

Schipani, E., G. S. Jensen, et al. (1997). "Constitutive activation of the cyclic adenosine 3',5'-monophosphate signaling pathway by parathyroid hormone (PTH)/PTH-related peptide receptors mutated at the two loci for Jansen's metaphyseal chondrodyplasia." *Mol Endocrinol* 11(7): 851-8.

Shenker, A., L. Laue, et al. (1993). "A constitutively activating mutation of the luteinizing hormone receptor in familial male precocious puberty [see comments]." *Nature* 365(6447): 652-4.

Sommers, C. M., N. P. Martin, et al. (2000). "A limited spectrum of mutations causes constitutive activation of the yeast alpha-factor receptor." *Biochemistry* 39(23): 6898-909.

Spalding, T. A., E. S. Burstein, et al. (1998). "Identification of a ligand-dependent switch within a muscarinic receptor." *J Biol Chem* 273(34): 21563-8.

Spalding, T. A., E. S. Burstein, et al. (1997). "Constitutive activation of the m5 muscarinic receptor by a series of mutations at the extracellular end of transmembrane 6." *Biochemistry* 36(33): 10109-16.

Tseng, C. C. and L. Lin (1997). "A point mutation in the glucose-dependent insulinotropic peptide receptor confers constitutive activity." *Biochem Biophys Res Commun* 232(1): 96-100.

Wonerow, P., T. Schoneberg, et al. (1998). "Deletions in the third intracellular loop of the thyrotropin receptor. A new mechanism for constitutive activation." *J Biol Chem* 273(14): 7900-5.

Light Emission Induced by the WT CCK-BR vs. a Constitutively Active Mutant




Figure 2

A Point Mutation Confers Constitutive Activity to the Rat μ Opiod Receptor




Figure 3

Forskolin Stimulated HEK293 Cells Transfected With pcDNA1 and a CRE-luc Construct




Figure 4

The Rat μ Opioid Receptor Signals Through G α i




Figure 5

A Point Mutation Confers Constitutive Activity to the Rat μ Opioid Receptor



Figure 6

Target Residues Within Class I GPCRs





Figure 7


TMD III Asn (-14 from DRY) is a Target for Mutation Induced Constitutive Activity



mu opioid
bradykinin B2
angiotensin II AT1A

Figure 8

The 'DRY' Motif is a Target for Mutation Induced Constitutive Activity



oxytocin
vasopressin-V2
cholecystokinin-A
melanocortin-4
 α_{1B} adrenergic

Figure 9

Figure 10. *WT* *MC-4* *D146M*

A Point Mutation Enhances MC-4 Receptor Constitutive Activity




Figure 10

The -13 Position is a Target for Mutation Induced Constitutive Activity




Figure 11

crk	1	-----MESPIQFRGEPCPTCAPSACI	PPNSSAWFPWGAEPI	DSNGSAGSEDIAQ
crkr	1	-----MESPIQFRGEPCPTCAPSACI	PPNSSAWFPWGAEPI	DSNGSAGSEDIAQ
orm	1	MDSSAAPTNASNCTDAEAYSSCSPAPS PGWVY	NLSHLDGQLSDPCGPNRTDLCGRDSL	
ormr	1	MDSSTGPGNTSDCSDPDAQASCSDA	PGSWI	NLSHVDCGQSDPCGLNRTGLCGNDSL
ord	1	-----MEFAPSAGABE	PPPLF	ASDAYPSACPSAGANASG
AT1a	1			MALNSSAEDGIKRI
BK-2	1	-----MFSPWKISMFLSVREDSVPTTASFS	ADMNVTLQGPTLNG	TFAD

crk	49	LEPAHISPAH..PMEHTAIVSVEVVGLAGNSLIVMBVIRYTKMKTATNLYIFNLALADA
orkr	49	LEPAHISPAH..PMEHTAIVSVEVVGLAGNSLIVMBVIRYTKMKTATNLYIFNLALADA
orm	59	CPPTGTS.PSMVTAITMIALYSHVCVVGLFGMFPLYNIVIRYTKMKTATNLYIFNLALADA
ormr	57	CPQTGTS.PSMVTAITMIALYSHVCVVGLFGMFPLYNIVIRYTKMKTATNLYIFNLALADA
ord	37	PPGARSASSIALAKAATTAIVSACAVGLAGNIVLWEGIRYTKMKTATNLYIFNLALADA
AT1a	16	DDCPHAGRHSYIFWVWPTDYSF:FWVGBFGNSLIVIVIYFVMGKIKIVASVELNLALADL
BK-2	45	SKCPQVEWLGLWLNTPQPPFLWVVEVATEENIFVLSVFCILHKSSCTVAELVYGNLAADL

ork	107	ENVVHTTPFQSTVYLMN . SWPPGIVLCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK
orkr	107	ENVVHTTPFQSAVYLMN . SWPPGIVLCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK
orm	118	LAISITLPEQSYNYLMG . SWPPGIVLCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK
ormr	116	LAISITLPEQSYNYLMG . SWPPGIVLCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK
ord	97	LAISITLPEQSAKYLME . SWPPGIVLCKAVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK
AT1a	76	CFLLTLELWAVYTAMEYRWRPFCGNHLCKIASASVTEMVYASMELLCTSEDRYIAVCHPVK
BK-2	105	ILACGILPEWATISNNFDWLREGETLQRVAVNATISMNLYSSICFLMVSSEDRYIAVCHPVK

ork	166	ALDFRTPLKAKLNICIWI SSVGISAIVKGTKVR..EDVDVIECSLQFDDDSWMD
orkr	166	ALDFRTPLKAKLNICIWI ASSVGISAIVKGTKVR..EDVDVIECSLQFDDDSWMD
orm	177	ALDFRTPLNAKLCINICW SSAIGPAMFMATIKYR..Q..GSIDCILTESHPTW.YWD
ormr	175	ALDFRTPLNAKLCINICW SSAIGPAMFMATIKYR..Q..GSIDCILTESHPTW.YWD
ord	156	ALDFRTPAKAKLNICIWI ASVGIVPVMVMAVTSPR..D..GAIVVOMLOFSPSW.YWD
AT1a	136	SRLRTMLVAKNTCIIWI MA GLASIVAEVILHRNV..YFIENTNITVCAFHYESRN.STLP
BK-2	165	MGRMRGVRWAKEYSIVIWCGLLTSSEPMVFRTMKEYSDEGHNVTA C SYPS..LIWE

ork	224	LFMKICVFIFAFMVPVLLIIVCYTLMLRLKSVRILSGSKEKDRNLRRITRIVLVVVAVF
orkr	224	LFMKICVFIFAFMVPVLLIIVCYTLMLRLKSVRILSGSKEKDRNLRRITRIVLVVVAVF
orm	232	NLKKICVFIFAFMVPVLLIIVCYGLMLRLKSVRILSGSKEKDRNLRRITRIVLVVVAVF
ormr	230	NLKKICVFIFAFMVPVLLIIVCYGLMLRLKSVRILSGSKEKDRNLRRITRIVLVVVAVF
ord	211	TVTKICVFIFAFMVPVLLIIVCYGLMLRLKSVRILSGSKEKDRSLRRITRIVLVVVAVF
AT1a	193	IGLGPDKNLLGFLFPFLILTSYIILWALKKKAYEIQKNKPRNDD...IFREDIMAVLFF
BK-2	222	VFTNVLLENVVGFLIP. LSWITFCIOMIIMQVLRNNEMQKFKEIQT. RRAVIVLVLVLLIF

ork	284	LVCWTPLHIFELVIALGS.T.....SHSTAAALSIVFCLALGYTNSSLNPVLYAFLDENF
orkr	284	ILVCWTPLHIFELVIALGS.T.....SHSTAAVLSSIVFCLALGYTNSSLNPVLYAFLDENF
orm	292	LVCWTPLHIFELVIALGS.T.....EIIIFQTVSWHEFCLALGYTNSSLNPVLYAFLDENF
ormr	290	LVCWTPLHIFELVIALGS.T.....EIIIFQTVSWHEFCLALGYTNSSLNPVLYAFLDENF
ord	271	LVCWTPLHIFELVWTLLDID.....RRDPLVVAALHLCLALGYANSSLNPVLYAFLDENF
AT1a	250	FFPSWVPHQIETPLVLLIQLGVIHDCKISDIDVDTAMPITCLAYFNNCLNPFLFYGLGKE
BK-2	280	ILVCWLPLFOISTPLTLHRIIGILSSCQDERIIDVITQIASFMAYNSNSCLNPVLYVIVGKRE

ork	338	KRCFRDFCFPLKMRMVEROSLSRMR. MIVMOD. PAYLRDIDGVMKPV-----	76
orkr	338	KRCFRDFCFPLKMRMVEROSLSRMR. MIVMOD. PASMRDVGGVMKPV-----	77
orm	346	KRCFRFECIPTSSNTEQONSTRFRONT. RDHPSTANTVDRTMHQLENLEAETAPLP	78
ormr	344	KRCFRFECIPTSSNTEQONSTRFRONT. RDHPSTANTVDRTMHQLENLEAETAPLP	79
ord	326	KRCFRQOLRKPCGRDPDPSFSRAREAAARERVTACTPSDGPGGGAAA-----	80
AT1a	310	KRYFLQLLKYIIPPKAKSHS...SLSTM..STLSYRPSDNMSSSAKKPASCPEVE-	81
BK-2	340	RKKSWEVYOGWCORGCRSEPIQMENSM..GTL..RTSISVERQIHKLQDWAGSRQ	82

SEQ ID NO:

Figure 12

mORMouse 1 MDSSAGPGMISDCSDP1A. PASCSPA. PGSWHNLSHADGNSDPOGPNTGGLGSHSLO
 mORrat 1 MDSSSTGPQNTSDCSDP1A. QASCSPA. PGSWHNLSHADGNSDPOGPNTGGLGNDNSLO
 mORbovin 1 MDSGAVPTMASNCDEPFTHPSSCSPAPSPSSWVAFSHLEGNLSDPOGPNTGGLGNDNSLO
 mORhuman 1 NDSSAAPTMASNCDEPFTHPSSCSPAPSPSSWVAFSHLEGNLSDPOGPNTGGLGNDNSLO
 mORpig 1 NDSSAAPTMASNCDEPFTSPSSMCSHPVPSPSSWVAFSHLEGNLSDPOGPNTGGLGNDNSLO
 mORws 1 MGS...GNIISDFLYPLS.....NIVMS....NSSVLCLRNFSNSTSFLNMNGSSRDSTD
 AT1a 1 ----- MALNSSAEDGKRIODDC
 BK-2 1 ----- MFSPWKISMFLSVREDSPPTTASFSADMNVTLQGETLNG. TFAQSKC

mORMouse 58 PGTGSPSMITAIIIMALYSIVCVGLPGMFLVMYIVIVRTKMKTAINTYIENLALADALA
 mORrat 58 PGTGSPSMITAIIIMALYSIVCVGLPGMFLVMYIVIVRTKMKTAINTYIENLALADALA
 mORbovin 61 HSAGSPSMITAIIIMALYSIVCVGLPGMFLVMYIVIVRTKMKTAINTYIENLALADALA
 mORhuman 60 PGTGSPSMITAIIIMALYSIVCVGLPGMFLVMYIVIVRTKMKTAINTYIENLALADALA
 mORpig 61 PGTGSPSMITAIIIMALYSIVCVGLPGMFLVMYIVIVRTKMKTAINTYIENLALADALA
 mORws 48 EODKIE. MIIAIITLYSIVCVGLPGMFLVMYIVIVRTKMKTAINTYIENLALADALA
 AT1a 19 EKAGRHSYLFVM. IPTLISIHFVVCHEGNSLIVVIVIYFYMKIKIVASVFLNLALADLCF
 BK-2 48 POVEWLGWPNITI. QPPFLWVIFVETLENIFVLSVFCILHKSSCIVABLYICNLAAADLIL

mORMouse 118 TSTLPPFOSVNYLMG. TWPEGNLLCKIVISIDYYNMFTSIFTLCIMSVDRYIAVCHPVKAL
 mORrat 118 TSTLPPFOSVNYLMG. TWPEGTLLCKIVISIDYYNMFTSIFTLCIMSVDRYIAVCHPVKAL
 mORbovin 121 TSTLPPFOSVNYLMG. TWPEGTLLCKIVISIDYYNMFTSIFTLCIMSVDRYIAVCHPVKAL
 mORhuman 120 TSTLPPFOSVNYLMG. TWPEGTLLCKIVISIDYYNMFTSIFTLCIMSVDRYIAVCHPVKAL
 mORpig 121 TSTLPPFOSVNYLMG. TWPEGTLLCKIVISIDYYNMFTSIFTLCIMSVDRYIAVCHPVKAL
 mORws 107 TSTLPPFOSVNYLMG. TWPEGTLLCKIVISIDYYNMFTSIFTLCIMSVDRYIAVCHPVKAL
 AT1a 78 LLTLPWAVYTAMEYRWPFCNHLCKIASASVTSVNLVYASVLLTCEHSDRYIAVHPMKS
 BK-2 107 ACGLPEWATITISNNFDWLFGETLCLRWVNAIIISMANLYSSTICFLMISIDRYIALVKTMSMG

mORMouse 177 DFRTPRANKIINVCNWLSSAIGLPVFMATTKYRQ.....GSIDCTLTFSHPTWYWE
 mORrat 177 DFRTPRANKIINVCNWLSSAIGLPVFMATTKYRQ.....GSIDCTLTFSHPTWYWE
 mORbovin 180 DFRTPRANKIINVCNWLSSAIGLPVFMATTKYRQ.....GSIDCTLTFSHPTWYWE
 mORhuman 179 DFRTPRANKIINVCNWLSSAIGLPVFMATTKYRQ.....GSIDCTLTFSHPTWYWE
 mORpig 180 DFRTPRANKIINVCNWLSSAIGLPVFMATTKYRQ.....GSIDCALTFSHPTWYWE
 mORws 166 DFRTPRANKIINVCNWLSSAIGLPVFMATTKYRQ.....GSIDCALTFSHPTWYWE
 AT1a 138 LFRIMLVAKUTCIIITWAGLASLPAVIRNV....YFIENTNITVCAHYESRNSTLP
 BK-2 167 RMRGVVERWAKLYSLVINGCILISSEPMIAVFRIMK...EYSDEGHNVTAQVISYPS..LIWE

mORMouse 230 NLLKICVPIFAPIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMLVUVVAVF
 mORrat 230 NLLKICVPIFAPIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMLVUVVAVF
 mORbovin 233 NLLKICVPIFAPIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMLVUVVAVF
 mORhuman 232 NLLKICVPIFAPIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMLVUVVAVF
 mORpig 233 NLLKICVPIFAPIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMLVUVVAVF
 mORws 226 TLLKICVPIFAPIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMLVUVVAVF
 AT1a 193 IGLGLTKNLLGFFPFLIILTSMLHKAQKAYELOKNPKRND...IPTLIMAFVLF
 BK-2 222 VFTNMLIENWVGELIP. LSVITFCIYDNOVLRNNEOKFKEIOTE. RRAVIVLVLVLLLF

mORMouse 290 IVCWTPPIHYVIIKALITI.....PETTIFOTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mORrat 290 IVCWTPPIHYVIIKALITI.....PETTIFOTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mORbovin 293 IVCWTPPIHYVIIKALITI.....PETTIFOTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mORhuman 292 IVCWTPPIHYVIIKALITI.....PETTIFOTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mORpig 293 IVCWTPPIHYVIIKALITI.....PETTIFOTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mORws 286 IVCWTPPIHYVIIKALITI.....PNSLFOTVSWHFCIALGYTNSCLNPVLYAFLDENF
 AT1a 250 FFSWVPHQITPFDVLIQFVHDCKISDIVDTAPITICLAYFNNCLNPVLYCFLGKIF
 BK-2 280 IICNLPPFOISTFUDTLHRIIGLSSCODERIIDVITOIASFM4YSNSCLNPVLYVIVGKRF

mORMouse 344 KRCFREFO. IPTSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLP 83
 mORrat 344 KRCFREFO. IPTSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLP 79
 mORbovin 347 KRCFREFO. IPTSSTIEQONSTRIRONTRHPSTANTVDRTNHOLENLEAETAPLP 84
 mORhuman 346 KRCFREFO. IPTSSTIEQONSTRIRONTRHPSTANTVDRTNHOLENLEAETAPLP 85
 mORpig 347 KRCFREFO. IPTSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLP 86
 mORws 340 KRCFREFO. IPTSSTIEQONSTRIRONTRHPSTANTVDRTNHOLENLEAETAPLP 87
 AT1a 310 KRYFLOLLKYLPKAKSHS...SLSTKMTLSYRPSDNSSSAKKPASCFEVE--- 81
 BK-2 340 RKKSWEVYOGUCOKGGCRSEPIQMENSMGTL..RISISVEROIKLQDNWASRQ--- 82

SEQ ID NO:

Figure 13